













Part. & Energy conservation [Case: I.1.5.j, Solver: 10, $D = 0.1 \text{ m}^2/\text{s}$, $v = -1.00 \text{ m/s}$, $\Delta t = 4.00$, $\tau = 1.0 \times 10^{-3} \text{ s}$, $N_p = 101$]
Comparison with initial solution - log scale; total time and zoom over time







Part. & Energy conservation [Case: I.1.5.j, Solver: 10, $D = 0.1 \text{ m}^2/\text{s}$, $v = -1.00 \text{ m/s}$, $\Delta t = 4.00$, $\tau = 1.0 \times 10^{-3} \text{ s}$, $N_p = 101$]
Comparison with initial solution - linear scale; total time and zoom over time



Part. & Energy conservation [Case: I.1.5.j, Solver: 10, $D = 0.1 \text{ m}^2/\text{s}$, $v = -1.00 \text{ m/s}$, $\Delta t = 4.00$, $\tau = 1.0 \times 10^{-3} \text{ s}$, $N_p = 101$]
Comparison with previous time-sampled (τ_{out}) solution - log and linear scales







Particle conservation [Case: I.1.5.j, Solver: 10, $D = 0.1 \text{ m}^2/\text{s}$, $v = -1.00 \text{ m/s}$, $\Delta t = 4.00$, $\tau = 1.0 \times 10^{-3} \text{ s}$, $N_p = 101$]
 Comparison with asymptotic solution (electrons and ions); total time and zoom over time







Profiles [Case: I.1.5.j, Solver: 10, $D = 0.1 \text{ m}^2/\text{s}$, $v = -1.00 \text{ m/s}$, $\Delta t = 4.00$, $\tau = 1.0 \times 10^{-3} \text{ s}$, $N_\rho = 101$]

Time sampling: total simulation time/10







Profiles [Case: I.1.5.j, Solver: 10, $D = 0.1 \text{ m}^2/\text{s}$, $v = -1.00 \text{ m/s}$, $\Delta t = 4.00$, $\tau = 1.0 \times 10^{-3} \text{ s}$, $N_\rho = 101$]

Comparison with asymptotic solution







Profiles [Case: I.1.5.j, Solver: 10, $D = 0.1 \text{ m}^2/\text{s}$, $v = -1.00 \text{ m/s}$, $\Delta t = 4.00$, $\tau = 1.0 \times 10^{-3} \text{ s}$, $N_\rho = 101$]

Time sampling: first 10 time slices or zoom over time $0.1 \times (a^2/D)/|1 - (Va/D)| = 0.19 \text{ s}$







0.00
0.04
0.08
0.12
0.16





Profiles [Case: I.1.5.j, Solver: 10, $D = 0.1 \text{ m}^2/\text{s}$, $v = -1.00 \text{ m/s}$, $\Delta t = 4.00$, $\tau = 1.0 \times 10^{-3} \text{ s}$, $N_\rho = 101$]
 Spatial zoom over magnetic axis; time sampling: first 10 time slices or zoom over time $0.1 \times (a^2/D)/|1 - (Va/D)| = 0.19 \text{ s}$



Profiles [Case: I.1.5.j, Solver: 10, $D = 0.1 \text{ m}^2/\text{s}$, $v = -1.00 \text{ m/s}$, $\Delta t = 4.00$, $\tau = 1.0 \times 10^{-3} \text{ s}$, $N_\rho = 101$]
 Spatial zoom over edge; time sampling: first 10 time slices or zoom over time $0.1 \times (a^2/D)/|1 - (Va/D)| = 0.19 \text{ s}$




Profiles [Case: I.1.5.j, Solver: 10, $D = 0.1 \text{ m}^2/\text{s}$, $v = -1.00 \text{ m/s}$, $\Delta t = 4.00$, $\tau = 1.0 \times 10^{-3} \text{ s}$, $N_\rho = 101$]
 Time sampling: last 10 time slices



Legend for time slices: 0.00, 0.04, 0.08, 0.12, 0.16, 0.20, 0.24, 0.28, 0.32, 0.36

Profiles [Case: I.1.5.j, Solver: 10, $D = 0.1 \text{ m}^2/\text{s}$, $v = -1.00 \text{ m/s}$, $\Delta t = 4.00$, $\tau = 1.0 \times 10^{-3} \text{ s}$, $N_\rho = 101$]
 Spatial zoom over magnetic axis; time sampling: last 10 time slices



Profiles [Case: 1.1.5.j, Solver: 10, $D = 0.1 \text{ m}^2/\text{s}$, $v = -1.00 \text{ m/s}$, $\Delta t = 4.00$, $\tau = 1.0 \times 10^{-3} \text{ s}$, $N_\rho = 101$]
 Spatial zoom over edge; time sampling: last 10 time slices

