Part. & Energy conservation [Case: I.1.5.j, Solver: 7, $D = 0.1\,\mathrm{m}^2/\mathrm{s}$, $v = -1.00\,\mathrm{m/s}$, Δt = 4.01, $\tau = 1.0 \times 10^{-2}\,\mathrm{s}$, $N_\rho = 101$]

Comparison with initial solution - log scale; total time and zoom over time









Part. & Energy conservation [Case: I.1.5.j, Solver: 7, $D = 0.1\,\mathrm{m}^2/\mathrm{s}$, $v = -1.00\,\mathrm{m}/\mathrm{s}$, Δt = 4.01, $\tau = 1.0 \times 10^{-2}\,\mathrm{s}$, $N_\rho = 101$]

Comparison with initial solution - linear scale; total time and zoom over time









Part. & Energy conservation [Case: I.1.5.j, Solver: 7, $D = 0.1\,\mathrm{m^2/s}$, $v = -1.00\,\mathrm{m/s}$, Δt = 4.01, $\tau = 1.0 \times 10^{-2}\,\mathrm{s}$, $N_\rho = 101$]

Comparison with previous time-sampled (τ_{out}) solution - log and linear scales

Particle conservation (electrons and ions)



Energy conservation (electrons and ions)







Particle conservation [Case: I.1.5.j, Solver: 7, $D = 0.1\,\mathrm{m}^2/\mathrm{s}$, $v = -1.00\,\mathrm{m}/\mathrm{s}$, Δt = 4.01, $\tau = 1.0 \times 10^{-2}\,\mathrm{s}$, $N_\rho = 101$]

Comparison with asymptotic solution (electrons and ions); total time and zoom over time









Profiles [Case: I.1.5.j, Solver: 7, $D = 0.1\,\mathrm{m}^2/\mathrm{s}$, $v = -1.00\,\mathrm{m/s}$, Δt = 4.01, $\tau = 1.0 \times 10^{-2}\,\mathrm{s}$, $N_\rho = 101$]

Time sampling: total simulation time/10









0.00
0.04
0.08
0.12
0.16
0.20
0.24
0.28
0.32
0.36
0.40
0.44
0.48
0.52
0.56
0.60

0.00
0.04
0.08
0.12
0.16
0.20
0.24
0.28
0.32
0.36
0.40
0.44
0.48
0.52
0.56
0.60

Profiles [Case: I.1.5.j, Solver: 7, $D = 0.1\,\mathrm{m^2/s}$, $v = -1.00\,\mathrm{m/s}$, Δt = 4.01, $\tau = 1.0 \times 10^{-2}\,\mathrm{s}$, $N_\rho = 101$]

Comparison with asymptotic solution









Profiles [Case: I.1.5.j, Solver: 7, $D = 0.1\,\mathrm{m}^2/\mathrm{s}$, $v = -1.00\,\mathrm{m}/\mathrm{s}$, Δt = 4.01, $\tau = 1.0 \times 10^{-2}\,\mathrm{s}$, $N_\rho = 101$]

Time sampling: first 10 time slices or zoom over time $0.1 \times (a^2/D)/|1 - (Va/D)|$ =0.19 s









Profiles [Case: I.1.5.j, Solver: 7, $D = 0.1\,\mathrm{m}^2/\mathrm{s}$, $v = -1.00\,\mathrm{m}/\mathrm{s}$, Δt = 4.01, $\tau = 1.0 \times 10^{-2}\,\mathrm{s}$, $N_\rho = 101$]

Spatial zoom over magnetic axis; time sampling: first 10 time slices or zoom over time $0.1 \times (a^2/D)/|1 - (Va/D)|$ =0.19 s









Profiles [Case: I.1.5.j, Solver: 7, $D = 0.1\,\mathrm{m}^2/\mathrm{s}$, $v = -1.00\,\mathrm{m/s}$, Δt = 4.01, $\tau = 1.0 \times 10^{-2}\,\mathrm{s}$, $N_\rho = 101$]

Spatial zoom over edge; time sampling: first 10 time slices or zoom over time $0.1 \times (a^2/D)/|1 - (Va/D)|$ =0.19 s









Profiles [Case: I.1.5.j, Solver: 7, $D = 0.1\,\mathrm{m}^2/\mathrm{s}$, $v = -1.00\,\mathrm{m/s}$, Δt = 4.01, $\tau = 1.0 \times 10^{-2}\,\mathrm{s}$, $N_\rho = 101$]

Time sampling: last 10 time slices









Profiles [Case: I.1.5.j, Solver: 7, $D = 0.1\,\mathrm{m}^2/\mathrm{s}$, $v = -1.00\,\mathrm{m/s}$, Δt = 4.01, $\tau = 1.0 \times 10^{-2}\,\mathrm{s}$, $N_\rho = 101$]

Spatial zoom over magnetic axis; time sampling: last 10 time slices









Profiles [Case: I.1.5.j, Solver: 7, $D = 0.1\,\mathrm{m}^2/\mathrm{s}$, $v = -1.00\,\mathrm{m/s}$, Δt = 4.01, $\tau = 1.0 \times 10^{-2}\,\mathrm{s}$, $N_\rho = 101$]

Spatial zoom over edge; time sampling: last 10 time slices